

KYRIOS clinical trial: Tracking the immune response to SARS-CoV-2 mRNA vaccines in an open-label multicenter study in participants with relapsing multiple sclerosis treated with ofatumumab s.c.

Tjalf Ziemssen¹, Tobias Bopp², Benjamin Etle³, Marie Groth³

¹Center of Clinical Neuroscience, Dresden University of Technology, Dresden, Germany.

²Institute for Immunology, University Medical Center, Mainz, Germany.

³Novartis Pharma GmbH, Nuremberg, Germany.

Session name: P1 - Poster Session 1

Session time: Apr 24 2022 9:00 AM - Apr 26 2022 5:00 PM

Poster Presentation at the American Academy of Neurology (AAN) 2022



Scan to download a
copy of this presentation

Tjalf Ziemssen has received research support, consulting fee, and honoraria for lectures from Alexion, Biogen, Celgene, Merck, Novartis, Roche, Sanofi, Teva.

Tobias Bopp has received consulting fee and honoraria for lectures from Biogen, Celgene, Merck, Novartis, Pathios Therapeutics, Roche, Sanofi Genzyme, Teva.

Benjamin Ettle and **Marie Groth** are employees of the Novartis Pharma GmbH, Nuremberg, Germany.

Funding source: This study was sponsored by Novartis Pharma Vertriebs GmbH.

- Development of SARS-CoV-2 vaccines was a key milestone in fighting the COVID-19 pandemic, but little is known about the efficacy of these vaccines in patients with Multiple Sclerosis (MS) treated with anti-CD20 therapies.
- Ofatumumab is the first fully-human anti-CD20 antibody authorized by the EMA for the treatment of adult patients with relapsing forms of multiple sclerosis (RMS) with active disease. Ofatumumab is applied once monthly s.c. (20 mg) and selectively depletes B-cells, which represent one pillar of the adaptive immune response.
- However, newly developed SARS-CoV-2 mRNA vaccines have been shown to not only induce selective B- but also T-cell responses^{1,2}.
- **Thus, it becomes essential to investigate both, humoral and cellular immune responses in patients treated with ofatumumab in order to provide guidance on vaccination for patients with MS and treating physicians.**

1. Sahin et al. (2021) Nature. 595,572–577. 2. Jackson et al. (2020) N Engl J Med. 383:1920-1931.

The aim of this study is to understand the impact of ofatumumab treatment on mounting cellular and humoral immune responses after initial and booster SARS-CoV-2 mRNA vaccination.

Background

Objective

Methods


Results

Conclusions



- KYRIOS is a two-cohort, open-label, prospective study currently including 33 RMS patients at eight sites in Germany (**Figure 1**).
 - This study compares the immune response in patients receiving SARS-CoV-2 mRNA vaccination prior to starting ofatumumab treatment (**cohort 1, control cohort**) and patients vaccinated during stable ofatumumab treatment for at least 4 weeks (**cohort 2**).
 - Immune responses after initial and booster vaccination were analyzed separately.
- Ofatumumab treatment is administered as part of the study and vaccinations as part of clinical routine according to summary of product characteristics (SmPC).
- **Neutralizing antibodies** were analyzed utilizing the cPassTMSARS-CoV-2 Neutralization Antibody Detection Kit from GenScriptUSA Inc(L00847).
- **SARS-CoV-2 specific T-cells** were detected with the CoV-iSpot Interferon- γ + Interleukin-2 (ELSP 7010 strip format) from GenID®GmbH. Each ELISpot assay was performed with 2×10^6 PBMCs (peripheral blood mononuclear cells).

Figure 1: Study design



Demographics and baseline information

- Patient characteristics at the time of screening are shown in **Table 1**.
 - There are currently 33 patients enrolled in the study with an average age of 41.6 years and a disease history of 6.7 years.
 - 50% in cohort 1 and 26% in cohort 2 were treatment naive.
- Most patients received BioNTech/Pfizer SARS-CoV-2 mRNA vaccines with an average of 4.8 weeks between 1st and 2nd dose.
- Booster vaccines were administered on average 6.1 months after 2nd dose and mostly (90%) with the same vaccine as in the initial vaccination cycle.
- B-cell depletion was verified in subjects of cohort 2 before vaccination.

Table 1: Patient characteristics


Variable*	Cohort 1 – vaccination prior to treatment	Cohort 2 – vaccination during stable treatment
N	14	19
Age, years	40.84 [23; 79]	42.08 [21; 61]
Sex, female, n (%)	10 (41.7)	11 (61.1)
Time since diagnosis, years	7.5 [0; 23]	6.1 [0; 19]
Prior treatments before ofatumumab		
Naive, N (%)	7 (50)	5 (26.3)
One, N (%)	2 (14.3)	4 (21.1)
Two, N (%)	0 (0)	5 (26.3)
More than two, N (%)	5 (35.7)	5 (26.3)
Vaccination, n (%)		
1 st (BioNTech Moderna)	13 (92.9) 1 (7.1)	18 (94.7) 1 (5.3)
2 nd (BioNTech Moderna)	13 (92.9) 1 (7.1)	18 (94.7) 1 (5.3)
Booster (BioNTech Moderna)	7 (87.5) 1 (12.5)	11 (84.6) 2 (15.4)
Vaccination time interval (days)		
1 st to 2 nd vaccination	30.8 [21; 42]	35.7 [21; 56]
2 nd vaccination to Booster	182 [160; 216]	187 [129; 295]
CD19+/CD20+ cells at baseline (cells/μl)	215.7 [7; 535]	0.1 [0; 1]

* if not indicated otherwise, data are presented as mean [min; max]; #depending on subcohort vaccine refers to either initial vaccination or booster vaccination

SARS-CoV-2 specific T-cell response

- All patients (5/5) receiving initial vaccination during stable ofatumumab treatment developed SARS-CoV-2 reactive T-cells (**Figure 2**).
- Extend of T-cell response in these patients was comparable to control cohort and peaked at 1 week after vaccination (data on file).
- T-cell response in boosted patients was comparable in both cohorts.
- Across both cohorts, most patients without T-cell response after booster (5/7) were older than 50 years.
- As this is an interim analysis, T-cell data after booster is still incomplete and needs to be interpreted with caution.


Figure 2: SARS-CoV-2 T-cell reactivity (IFN- γ and/or IL-2)



Development of SARS-CoV-2 neutralizing antibodies

- Neutralizing antibodies (NAb) represent only **a subset of all specific antibodies** and are considered a **more stringent** correlate of protective immunity. Total anti-SARS-CoV-2 IgGs were not measured here but might further contribute to immunity.
- All patients (4/4) receiving their initial vaccination during stable ofatumumab treatment had an increase in NAb (**Figure 3**).
- 50% of ofatumumab patients exceeded the assay-specific cut-off for seropositivity one week after vaccination.
- 5/6 patients boosted during stable ofatumumab treatment showed an increase in NAb to a similar extend as in control group.
- One patient who was seronegative before booster seroconverted during stable ofatumumab treatment.

Figure 3: Development of SARS-CoV-2 neutralizing antibodies



Safety

- One MS relapse occurred during the study (patient in cohort 2, relapse occurred before 1st vaccination, patient recovered fully)
- Two patients developed COVID-19 infections during the study:
 - One patient in cohort 1 (initial vaccination prior to ofatumumab treatment): 6 days after 2nd vaccination, CTCAE moderate, no MS therapy at time of infection, full recovery (duration of infection: 9 days)
 - One patient in cohort 2 (initial vaccination during ofatumumab treatment): 27 days after 2nd vaccination, CTCAE moderate, no interruption of ofatumumab treatment, infection was ongoing at time of data cut-off but has by now fully recovered (duration of infection: 13 days)

- **All patients (5/5) vaccinated during continuous ofatumumab treatment developed an immune response as soon as one week after initial vaccination cycle.**
 - T-cell response was not affected by ofatumumab treatment and similar to control cohort
 - All patients showed an increase in neutralizing antibodies. Although the extent was lower versus control group, 50% exceeded the cut-off value for seropositivity.
 - These results are in line with previously reported low rate of COVID-19 infections in vaccinated patients treated with ofatumumab³
- **Immune response after booster vaccine was similar in patients boosted before and during ofatumumab treatment.**
 - 6/7 ofatumumab patients had increased neutralizing antibodies to a comparable extend as in control cohort
 - one patient who was seronegative before booster seroconverted during continuous ofatumumab treatment
 - T-cell response was heterogenous but comparable between cohorts. However, T-cell data is still incomplete and needs to be interpreted with caution.
- Despite limited sample size, population heterogeneity and pending longitudinal data, we can conclude that both cellular and humoral response need to be considered for interpretation of vaccine efficacy.

3. Cross et al. (2022) *Neurology and Therapy*

